My Colonoscopy Prep Calendar—Bowel Prep using SUPREP (for procedures at BASC)













Your Pharmacy Shopping List:

- ☐ One bottle Milk of Magnesia (or generic equivalent of magnesium hydroxide).
- ☐ Prescription for the SUPREP.

Your Grocery Shopping List:

- ☐ Purchase clear liquid products— examples include:
 - sports drinks (not red, blue or purple). Diabetic patients are recommended to purchase a sugar-free version.
 - apple or white grape juice
 - ginger ale
 - clear broth or bouillon
 - water
 - coffee/tea without milk
 - clear carbonated beverages
 - plain Jell-O (not red, blue or purple and no fruit or toppings)
 - Italian ice

- ☐ STOP eating all fruits, vegetables, seeds and nuts.
- ☐ Stop taking any iron or fiber supplements.
- supplements.

 Confirm ride and contact phone number for your colonoscopy. You MUST have a ride home from a responsible adult as
- received sedation for the procedure. (**NO** taxis, Lyft, Uber, etc).

vou will have

☐ Please inform the person providing your ride that the appointment will take approximately 2 ½-3 hours.

- ☐ Only consume
 CLEAR LIQUIDS
 THE ENTIRE DAY
 (see 7 days list for suggested liquids).
- □ DO NOT eat any solid food.
- ☐ It is important that you drink lots of fluids 2 days before your procedure as the preparation is very dehydrating.
- ☐ Take 2 tablespoons of Milk of Magnesia at dinner time.

- ☐ Consume only
 CLEAR LIQUIDS the
 entire day (no solid
 food, dairy, juice with
 pulp, alcohol or
 anything red, blue or
 purple in color).
- ☐ In the morning:

 Prepare the solution of SUPREP and refrigerate.

4 PM:

☐ Drink one of the 6 oz. bottles of SUPREP and refrigerate.

- ☐ Have NOTHING to eat or drink after midnight, except for finishing the SUPREP and/or taking any medications as ordered.
- □ 5 hours before
 your check-in
 time: FINISH the
 SUPREP by
 drinking the second
 6 oz. bottle.

Important:

☐ If you are on

blood pressure

medication, you

must take your

medication the

morning of your

procedure, prior to

check-in.



Please see important medical information on the reverse side

If you are **unable** to make your appointment, please call: **603.665.2470**.

New Hampshire Gastroenterology requires **72 hour notice** for cancellations, as we maintain a waiting list for these appointments.

Thank you for your consideration.

My Colonoscopy Prep Calendar—Bowel Prep using SUPREP (for procedures at BASC)

IMPORTANT information

Please continue your daily medication regimen except for:		Please bring a list of your current medication to the procedure and include dosing information.
	Diabetic Patients: Do not take any diabetes medication on the day of your procedure. If you have any questions, please call your prescribing provider. If you take insulin, call your prescribing provider and ask for instructions.	You will be contacted by Endoscopy one weel prior to your procedure to review important details. You will also be contacted 48 hours prior to the procedure to be given your arrival time.
	Patients on Blood Thinners: YOU MUST check in with your prescribing physician before stopping your medication. If you are currently taking an ASPIRIN daily, do not stop. Below is a list of common blood thinners and when, with physician approval, to discontinue them:	If you have any discomfort around the perianal region, use wet wipes instead of toilet paper and apply Vaseline or A&D Ointment to the rectum.
	 Ticlid (ticlopidine)—stop 10 days prior to procedure Effient (prasugrel)—stop 7 days prior to procedure Coumadin (warfarin), Plavix (clopidogrel) or Brilinta—stop 5 days prior Pradaxa (dabigatron), Eliquis (apixaban) or Xarelto (rivaroxaban)—stop 48 hours prior 	If you are unable to make your appointment please call: 603.665.2470. New Hampshire Gastroenterology requires 72 hour notice for cancellations, as we maintain a waiting list for these appointments. Thank you for your consideration.
	Patients on Erectile Dysfuntion medication: Viagara, Cialis, Adcirca, Sildenafil or Revatio—stop 24 hours prior.	



My Procedure Preparation Instructions (for procedures at BASC)

Diabetic and Weight Loss (Appetite Suppressant) Medications

If you are taking the following medications, you must ensure that your procedure takes place at least 8 days after your most recent dose:

Generic Drug	Brand Name
Albiglutide	Tanzeum, Eperzan
Dulaglutide	Trulicity
Exenatide (ER)	Bydureon Bcise
Exenatide (IR)	Byetta
Liraglutide (3mg)	Saxenda
Liraglutide (1.2mg, 1.8mg)	Victoza
Lixisenatide	Adlyxin, Lyxumia
Naltrexone-Bupropion others	Contrave
Orlistat	Alli
Others	PhenQ
Phentermine, Phentermine/Topiramate	Lomaira, Adipex-P, Qsymia
Pramlintide	Symlin
Semaglutide	Ozempic, Wegovy, others
Semaglutide	Rybelsus
Setmelanotide	Imcivree
Tirzepatide	Mounjaro, Zepbound

 Please ensure you are checking your blood sugar levels at a more frequent rate while you are holding these medications.

