My Colonoscopy Prep Calendar—One Day Bowel Prep



Your Pharmacy

Shopping List:

7 days before procedure



7 days procedure

Your Grocery Shopping List:

- ☐ 5mg Dulcolax tablets ☐ Purchase clear (or generic equivalent called bisacodyl).
- ☐ One 238g (8.3oz) bottle of Miralax (or generic equivalent called polyethylene glycol).
- ☐ One 64oz Gatorade. Pedialyte or any clear liquid of your choice (Any color EXCEPT for RED. BLUE or PURPLE).
- ☐ One bottle Milk of Magnesia (or generic equivalent magnesium hydroxide).

- liquid products examples include:
 - sports drinks (not red, blue or purple)
 - apple or white grape juice
 - ginger ale
 - clear broth or bouillon
 - water
 - coffee/tea without milk
 - clear carbonated beverages
 - plain Jell-O (not red, blue or purple and no fruit or toppings)
 - Italian ice



5 days before procedure

- ☐ Stop taking any iron supplements.
- ☐ Confirm ride and contact phone number to your colonoscopy. You MUST have a ride home from a responsible adult as you will have received sedation for the procedure. (NO taxis, Lyft, Uber, etc).
- ☐ Please inform the person providing your ride that the appointment will take approximately $2\frac{1}{2}$ hours.



- 3 days before procedure
- ☐ STOP eating nuts, seeds, corn. beans or any vegetable with a skin.
- ☐ STOP taking fiber supplements.
 - If you are **unable** to make your appointment, please call: 603.665.2470.

New Hampshire Gastroenterology requires 72 hour notice for cancellations, as we maintain a waiting list for these appointments.

Thank you for your consideration.



2 days before procedure

☐ Take 2 tablespoons of Milk of Magnesia at dinner time.



1 dav procedure

- ☐ Consume only CLEAR LIQUIDS the entire day (no solid food, dairy, juice with pulp, alcohol or anything red, blue or purple in color).
- ☐ **3 PM**: Take 10 mg total of Dulcolax.
- ☐ 6 PM: Mix Miralax 238g bottle with 64oz. Gatorade and shake until dissolved. Drink 8oz every 10-15 minutes until the bottle is HALF EMPTY (drink it fast). It will take approximately one hour to drink half the solution. Expect some discomfort.



- ☐ Have **NOTHING** to eat or drink after midnight.
- □ 5 hours before your check-in time: FINISH the Miralax drinking 8oz. every 10-15 minutes until the bottle is empty

Important:

☐ If you are on **blood** pressure medication. you must take your medication the morning of your procedure, prior to check-in.





Please see important medical information on the reverse side

My Colonoscopy Prep Calendar—One Day Bowel Prep

IMPORTANT information

☐ Please bring a list of your current Please continue your daily medication regimen except medication to the procedure and include for: dosing information. **Diabetic Patients:** Do not take any diabetes ☐ You will be contacted by Endoscopy **one week** medication on the day of your procedure. If you **prior** to your procedure to review important have any questions, please call your prescribing details. You will also be contacted 48 hours provider. If you take insulin, call your prescribing prior to the procedure to be given your arrival provider and ask for instructions. time. ☐ Patients on Blood Thinners: YOU MUST check ☐ If you have any discomfort around the in with your prescribing physician before stopping perianal region, use wet wipes instead of toilet your medication. If you are currently taking an paper and apply Vaseline or A&D Ointment to ASPIRIN daily, do not stop. Below is a list of the rectum. common blood thinners and when, with physician approval, to discontinue them: ☐ If you are unable to make your appointment, ■ Ticlid (ticlopidine)—stop 10 days prior to please call: 603.665.2470. New Hampshire procedure Gastroenterology requires **72 hour notice** for Effient (prasugrel)—stop 7 days prior to procedure cancellations, as we maintain a waiting list for Coumadin (warfarin), Plavix (clopidogrel) or these appointments. Thank you for your Brilinta—stop 5 days prior consideration. Pradaxa (dabigatron), Eliquis (apixaban) or Xarelto (rivaroxaban)—stop 48 hours prior □ Patients on Erectile Dysfuntion medication: Viagara, Cialis, Adcirca, Sildenafil or Revatio—stop



24 hours prior.