Preparation for your Esophagogastroduodenoscopy (EGD)

IMPORTANT information

Please continue your daily medication regimen except for:	 Please bring a list of your current medication to the procedure and include 	
Diabetic Patients: Do not take any diabetes medication on the day of your procedure. If you have any questions	dosing information.	
please call your prescribing provider. If you take insulin, call your prescribing provider and ask for instructions.	You will be contacted by Endoscopy one week prior to your procedure to review important details. You will also be contacted	ed
Please see the following if you are having a dilation, ablation (Barrx, or endoscopic mucoscal resection (EMR):	48 hours prior to the procedure to be given your arrival time.	
 Patients on Blood Thinners: YOU MUST check in with your prescribing physician before stopping your medication. If you are currently taking an ASPIRIN daily do not stop. Below is a list of common blood thinners and when, with physician approval, to discontinue them: Ticlid (ticlopidine)—stop 10 days prior to procedure Effient (prasugrel)—stop 7 days prior to procedure Coumadin (warfarin), Plavix (clopidogrel) or Brilinta—stop 5 days prior 	☐ You will be fasting prior to your procedure Have nothing to eat or drink after MIDNIGHT the night before your procedure. If you have questions regarding which medications you should or should not take the morning of your procedure, pleas contact the prescribing provider's office. We medications can be postponed until after your procedure is completed.	ng ot se Iost
 Pradaxa (dabigatron), Eliquis (apixaban) or Xarelto (rivaroxaban)—stop 48 hours prior 	 If you are unable to make your appointment, please call: 603.665.2470. New Hampshire Gastroenterology require 	s 7 2
 Patients on Erectile Dysfuntion medication: Viagara, Cialis, Adcirca, Sildenafil or Revatio—stop 24 hours prior 	hour notice for cancellations, as we main a waiting list for these appointments. Than you for your consideration.	

